



USAID
FROM THE AMERICAN PEOPLE

DHS WORKING PAPERS

HIV Knowledge and Risky Sexual Behavior among Men in Rwanda

Etienne Rugigana
Francine Birungi
Manassé Nzayirambaho

2014 No. 105

September 2014

This document was produced for review by the United States Agency for International Development.

DEMOGRAPHIC
AND
HEALTH
SURVEYS

HIV Knowledge and Risky Sexual Behavior among Men in Rwanda

Etienne Rugigana¹

Francine Birungi¹

Manassé Nzayirambaho¹

ICF International

Rockville, Maryland, USA

September 2014

¹School of Public Health, University of Rwanda

Corresponding author: Manassé Nzayirambaho, School of Public Health, University of Rwanda, Kigali, Rwanda;
Email: mnzayira@nursph.org

ACKNOWLEDGMENTS

This working paper was funded by USAID through the DHS Fellows program implemented by ICF International. The authors are very grateful for comments, reviews, and suggestions made by Sarah Staveteig and Wang Wenjuan. Particularly, we are grateful to Sarah Staveteig and to Dana Thomson for the constant technical support and advice they provided to us. Special thanks to Joy Fishel for reviewing our paper and providing invaluable comments. Also we would like to extend our sincere appreciation to all the 2014 DHS Fellows and co-facilitators for their support throughout the Fellows Program. We are also very grateful to Bryant Robey for editing, Shireen Assaf for assistance checking numbers in the document and Yuan Cheng for formatting.

The *DHS Working Papers* series is an unreviewed prepublication series of papers reporting on research in progress that is based on Demographic and Health Surveys (DHS) data. This research is carried out with support provided by the United States Agency for International Development (USAID) through The DHS Program (#AID-OAA-C-13-00095). The views expressed are those of the authors and do not necessarily reflect the views of USAID or the United States Government.

The DHS Program assists countries worldwide in the collection and use of data to monitor and evaluate population, health, and nutrition programs. For additional information about the DHS Program contact: DHS Program, ICF International, 530 Gaither Road, Suite 500, Rockville, MD 20850, USA. Phone: +1 301-407-6500; Fax: +1 301-407-6501; Email: reports@dhsprogram.com; Internet: www.dhsprogram.com.

ABSTRACT

Introduction: In many countries different interventions have been implemented to prevent new infections of the Human Immunodeficiency Virus (HIV). Rwanda has developed and implemented many strategies at the national level to reduce the incidence of HIV in the general population. One of the main objectives of such interventions is to improve the general level of knowledge of HIV, with the hypothesis that increasing HIV knowledge will reduce risky sexual behavior. However, there has been a concern that HIV knowledge may not necessarily reduce risky sexual behavior. Only a limited number of population-based studies describe the results of these interventions in terms of how HIV knowledge affects risky sexual behavior. Therefore, the aim of this paper is to fill in this gap, by exploring HIV knowledge and its effect on risky sexual behavior among men in Rwanda.

Methods: The study used data from the 2005 and 2010 Rwanda Demographic and Health Surveys to analyze the association between HIV risk factors and two types of risky sexual behavior—having two or more partners in the past 12 months; and among those with two or more partners, not using a condom at last sex—and the association between HIV knowledge and those risky behaviors. In addition to descriptive and bivariate analyses, multivariate logistic regression on pooled data from 2005 and 2010 was used to determine predictors of risky sexual behavior.

Results: Among men in the study sample, 5% in 2005 and 7% in 2010 reported having two or more sexual partners in the last 12 months preceding the survey. Among these men, 93% in 2005 and 74% in 2010 did not use a condom at the last sex. Knowledge of the protective effect of having just one uninfected faithful partner decreased from 90% to 82% between 2005 and 2010, and basic knowledge of HIV decreased from 71% in 2005 to 68% in 2010. Knowledge of the protective effect of using condoms increased from 90% in 2005 to 94% in 2010. In adjusted multivariate analysis, HIV knowledge was not associated with either type of risky sexual behavior.

Conclusion: The findings suggest that, while knowledge of the protective effect of using condoms has increased, knowledge of the protective effect of one uninfected faithful partner has been decreasing in Rwanda. This should draw attention to the need to take corrective measures. The study did not find strong evidence of a significant association between HIV knowledge and

risky sexual behavior among men in Rwanda. In setting up policies and strategies related to HIV prevention, policymakers should consider that focusing on HIV knowledge is not sufficient in itself.

Keywords: HIV knowledge, men, risky sexual behavior, Rwanda

INTRODUCTION

In Rwanda as in many other countries, different interventions have been implemented to fight against new infections of the Human Immunodeficiency Virus (HIV). Generally, one of the main objectives of such interventions is to improve general knowledge of HIV. Around the world, local and international efforts have led to successful HIV prevention programs in terms of improving HIV knowledge and awareness. However, HIV transmission is not decreasing as rapidly as desired, especially when considering the large amount of money and many other efforts spent on its prevention (UNAIDS 2008).

It is therefore important to understand the relationship between HIV knowledge and risky sexual behavior (RSB), and the different components of HIV knowledge that are positively associated with the reduction of RSB.

Since 1987, the government of Rwanda has been making substantial efforts in the fight against HIV/AIDS. Activities related to HIV/AIDS prevention and treatment have been carried out through different structures such as the National Program for the Fight against AIDS (PNLS), the National AIDS Commission (CNLS), the Treatment and Research AIDS Centre Plus (TRAC Plus), and the district AIDS committees (CDLS). Different strategies used in the fight against AIDS are also found in the Rwandan National AIDS strategic Plan. These include Information, Education, and Communication (IEC) strategies, which target mainly sexually active men age 15-59 and women age 15-49 (MOH [Rwanda] and RBC 2009; Hong et al. 2013).

These efforts have resulted in the achievement of two objectives: first, increased awareness of the existence of HIV as well as its transmission. Data indicate that the proportion of men who have heard of HIV/AIDS was virtually universal, both in 2005 and 2010 (NISR et al. 2012). The second objective is increased awareness of ways to prevent HIV infection. Under this objective, information related to the “ABC” strategy (abstinence, being faithful, using a condom) has been widely disseminated. Condoms have also been distributed free of charge in different strategic places or sold at very low cost, from the cities to the last small shop in villages.

Despite a satisfactory situation in terms of knowledge of HIV prevention and availability of condoms, many people still engage in RSB.

Background

RSB can be defined in different ways. The U.S. Centers for Disease Control and Prevention (CDC) defines RSB as behavior that increases one's risk of contracting sexually transmitted infections and experiencing unintended pregnancies. Risky behaviors are measured through a number of elements that include having sex at an early age, having multiple sexual partners, having sex while under the influence of alcohol or drugs, and having unprotected sex (CDC 2010). Many authors include in their definitions of RSB two main elements: 1) having sexual intercourse with a casual acquaintance without using a condom; 2) having multiple sexual partners (Booyesen 2004).

This study is about HIV-related RSB. We focus on two types of RSB among men that have been identified by international experts as indicators of risky behavior: multiple sexual partners (having two or more sexual partners in the past 12 months) and unprotected sex (not using a condom at the last sex for someone who had two or more sexual partners in the past 12 months). These two indicators—P8.11.N and P8.12.N, respectively, in the President's Emergency Plan for AIDS Relief (PEPFAR)¹—are useful measures of risk behavior in the population (PEPFAR 2013).

Different studies have reported low levels of condom use regardless of the infection risks (de Walque and Kline 2009; Biraro et al. 2009; Ahmed et al. 2001), while some others did not find significantly higher levels of condom use among populations with higher prevalence of HIV (Lagarde et al. 2001). A study conducted in Ethiopia among factory workers with a high HIV prevalence also showed low condom use despite widespread of knowledge of condoms (Sahlu et al. 1999). Similarly, an increase of some types of RSB was observed in Uganda over the period of 2001-2005, where a higher percentage of males age 15–49 in 2005 reported having multiple sexual partners and having sex with non-spousal partners compared with 2001 (25% versus 29% and 28% versus 37%, respectively) (Opio et al. 2008). Such a trend in RSB is notable given the high level of HIV prevention knowledge and known reduced risk of contracting HIV in the case of consistent condom use (Weller and Davis 2002; Pinkerton and Abramson 1997). This is why the question arises about the effect of HIV knowledge on RSB.

¹ PEPFAR defines P8.11.N and P8.12.N among women and men age 15-49; this report uses men age 15-59.

The literature gives mixed results about the effect of HIV knowledge on RSB. For example, studies in Zambia, Uganda, and Kenya show a positive effect of knowledge of HIV prevention methods on protective sexual behavior (MacIntyre et al. 2001; Magnani et al. 2002). However, other studies reveal little impact on condom use of having good knowledge of its protective benefits (Essien et al. 2002; Williams et al. 2003). Empirical theories about the link between behavior and knowledge suggest that self-perceived vulnerability to HIV is probably the main factor underlying RSB, more so than knowledge (McKusick and Coates 1985). The basis of HIV perceived vulnerability is people's beliefs about their own behavior as well as the possibility of getting HIV from this behavior (Gillespie 1997). Perceived vulnerability is an important component of the Health Belief Model. Specifically for HIV, it has been proposed that in the general population there is a negative association between RSB and HIV self-perceived vulnerability (McKusick and Coates 1985). That is, when people perceive that they are less vulnerable to HIV, they would be more likely to engage in RSB, independent of their knowledge about HIV.

In addition to HIV knowledge, some studies have also explored other factors correlated with RSB. According to the literature of developed countries, associations between RSB and socio-demographic factors are fairly robust (Kumari and Nair 2012). In other studies, some individual characteristics such as age, religion, caste, wealth status, education, occupation, and knowledge about sexually transmitted infections and HIV were associated with increased RSB (Kimuna and Djamba 2005; Li et al. 2009; Gras et al. 2001; Seidman et al. 1994). Other factors associated with higher RSB are smoking and alcohol (Opio et al. 2008; Weller and Davis 2002; Pinkerton and Abramson 1997), as well as witnessing violence among parents (Pinkerton and Abramson 1997). Recent studies also have reported an association between male circumcision and RSB (Kibira et al. 2014; Cassell et al. 2006; Eaton and Kalichman 2009).

To sum up, many strategies aiming at reducing RSB assume that increasing HIV knowledge and making condoms available will increase safer sexual behavior. However, it is not always the case. The present study assesses this relationship between HIV knowledge and RSB in the context of Rwanda, where programmatic efforts in HIV prevention measures have been scaled up between 2005 and 2010.

Research Questions

In Rwanda, HIV new infections are estimated to be 6,000 per year (MOH [Rwanda] and RBC 2014). HIV prevalence is 3% in the general population, and 2.2% in men versus 3.7% in women. This prevalence is highest among women in urban areas, at 8.7%. It was also found that HIV prevalence is highest among those who are widowed, divorced, or separated, and those who are in polygamous unions (MOH [Rwanda] and RBC 2014).

Even though men's HIV prevalence in Rwanda is relatively low compared with some other African countries, Rwanda ranks 13th among 27 sub-Saharan African countries (ICF International 2012), unchanged since 2005. Thus HIV still constitutes a big burden on the Rwanda Health System. Many strategies have been developed and implemented at the national level to reduce the incidence of HIV in the general population of Rwanda. These strategies include mass media communication, community events and outreach, and campaigns. Young people have been reached through school-based sexual health and AIDS education, anti-AIDS clubs, and outreach work for out-of-school youth. One of the main objectives has been to provide the entire population with comprehensive knowledge of HIV/AIDS (MOH [Rwanda] and RBC 2009).

However, very few population-based studies describe the results of these interventions in terms of status of HIV knowledge and its effect on RSB. Thus, the aim of this study is to fill in this gap, exploring HIV knowledge and its effect on RSB among men, based on data from the 2005 and 2010 Rwanda Demographic and Health Surveys (RDHS). Specifically, this study tries to answer the following questions:


- Are there differences in HIV knowledge between the 2005 and 2010 surveys?
- What are the differences in RSB between the 2005 and 2010 surveys?
- What is the association between HIV knowledge and RSB of men in 2005 and 2010?

The answer to these questions can help to design more focused and strategic interventions. This is particularly necessary not only for Rwanda but also for other comparable countries where governments and partners continue to implement and scale up HIV prevention interventions. It is important to implement HIV prevention programs based on strong evidence. This implies better understanding of the HIV dynamics and determinants that drive the

transmission of new infections, and thus selecting and focusing on only those interventions that effectively and efficiently contribute to preventing new infections.

Figure 1 shows the assumed factors in predicting engagement of men in RSB. Background characteristics (age, occupation, education, marital status, religion, residence, and wealth) constitute the risk factors of engaging in RSB [having two or more sexual partners in the past 12 months, or not using a condom at the last sex for someone who had two or more sexual partners in the past 12 months]. The framework postulates also that knowledge of HIV can modify the effect of risk factors on RSB, which means that men who have the above-mentioned risk factors should engage in fewer risky behaviors if at the same time they have knowledge of HIV.

Figure 1. Factors model for predicting risky sexual behavior among men



DATA AND METHODS

Data


Data from the 2005 and 2010 Rwanda Demographic and Health Surveys (RDHS) (NISR et al. 2006; NISR et al. 2012) have been used to analyze, on one hand, the association between risk factors and RSB, and on the other hand, the association between knowledge of HIV and RSB. The National Institute of Statistics of Rwanda and the Ministry of Health implemented the two RDHS surveys. ICF International provided technical assistance for the surveys through the USAID-funded MEASURE DHS program. The two RDHS samples were nationally representative, household-based surveys, designed to provide population and health indicator estimates at the national, urban-rural, and district levels.

The sampling method used was stratified two-stage cluster sampling. The first stage consisted of selecting clusters (enumeration areas) from each stratum (district). The second stage was a systematic sampling of households from each cluster selected. Then selected households were visited and interviewed. The eligible men's response rate was 97.2 % and 98.7% in 2005 and 2010 respectively. Eligible men age 15-59 were interviewed in half of the households. However, during analysis, only men who said that they had sexual intercourse during the last 12 months were considered, and this led our study population to 2 773 men in 2005 and 3 772 men in 2010 (figure 2). The two surveys were conducted from February to July (2005 RDH) and from September 2010 to March 2011 (NISR et al. 2012).

Key Variables and Measurements

During the interview, the following information on RSB was collected in the Men's Questionnaire: sexual activities in the 12 months preceding the survey, the number of sexual partners in the past 12 months (and the relationship with each partner for up to three partners), and condom use in the last sexual intercourse with each partner (for up to three partners).

Figure 2. Derivation of sample



In total, 4,820 men in 2005 and 6,329 men in 2010 were interviewed. Among them, 2,773 men in 2005 and 3,772 men in 2010 had sexual intercourse during the last 12 months. Among those who had sexual intercourse, 138 men in 2005 and 262 in 2010 had two or more sexual partners (Figure 2).

In this study, RSB was defined as having two or more sexual partners in the past 12 months, or not using condom at the last sex among individuals who had two or more sexual partners in the past 12 months. In fact, this is a metric of inconsistent condom use. The issue is that when a man has multiple partners and does not use condom consistently, he puts both partners at the risk of HIV infection, even if the most recent partner is a faithful spouse.

HIV knowledge was defined as knowing that consistent use of condom during sexual intercourse and having just one uninfected faithful partner can reduce the chance of getting the AIDS virus, as well as having “basic knowledge” about HIV (knowing that a healthy-looking person can have the AIDS virus, and rejecting the two following most common local misconceptions about AIDS transmission or prevention: HIV transmission by mosquito bites and sharing food).

Statistical Analysis

Analysis was performed using Stata 11 (Stata Corporation, College Station, Texas) to find the prevalence of RSB according to background characteristics (age, occupation, education, marital status, religion, residence, and wealth). Bivariate analysis was carried out and t-test was used to assess any significant change during the five-year study period. Also, logistic regression was done to determine predictors of RSB. Two binary outcomes variables were: “having two or more sexual partners in the past 12 months” and “not using condom at the last sex for someone who had two or more sexual partners in the past 12 months”. The following variables were tested in the model: knowledge of HIV and background characteristics. Chi-square test at the level of 5% was used to determine the statistical significance while associations were measured using odds ratios (OR) with 95% confidence intervals (CI). To account for the survey design and non-response, all analyses applied complex sampling weights.

RESULTS

A total of 2,773 men in 2005 and 3,772 men in 2010 reported that they had sex during the last 12 months preceding the survey. Among them, 138 in 2005 and 262 in 2010 declared that they had two or more sexual partners.

Sample Characteristics

As Table 1 shows, the age composition of sexually active men remained fairly stable between 2005 and 2010, with men age 15-24 comprising the largest share of that group in both years (43% and 41%, respectively). Among sexually active men, however, those with two or more partners tended to be age 25 or older, with 32% and 40% being in the 25-34 age group in 2005 and 2010, respectively. The majority of men who had two or more sexual partners had primary education level (75% in 2005 and 69% in 2010), were located in rural areas, and were married (83% in 2005 and 73% in 2010).

Among sexually active men, wealth index status remained stable between 2005 and 2010, with the middle, richer, and richest quintiles comprising the largest share. However, among sexually active men, those who had two or more partners tended to be in poorer (29%) and middle (24%) quintiles in 2005, compared with middle (25%) and richest (26%) in 2010.

The occupation profile of sexually active men varied from 2005 to 2010. Men who reported not working decreased substantially, from 36% in 2005 to just 1% in 2010, while jobs related to professionals/clerical/sales/services increased from 8% in 2005 to 43% in 2010. These changes are associated with the governance between the two surveys, especially the “Economic Development and Poverty Reduction Strategy” (EDPRS I) established in 2008. The EDPRS focused on job creation and the reduction of underemployment by making micro loans through microfinance cooperatives across the country.

Table 1 also shows that the majority of men with two or more sexual partners have the following main background characteristics: married, age 25-34, agriculture occupation, and rural location.

Table 1. Sample characteristics

	Among men who had sex last year		Among men who had 2+ partners last year	
	2005 (n=2,773)	2010 (n=3,772)	2005 (n=138)	2010 (n= 262)
Age (years)				
15-24	42.5	41.2	11.0	16.6
25-34	23.7	27.6	32.0	40.2
35-44	17.5	14.5	28.2	18.5
45-54	13.2	12.6	24.4	18.5
55-59	3.1	4.1	4.4	6.2
Education level				
No education	22.9	16.6	16.0	13.0
Primary	64.8	68.8	74.7	69.4
Secondary	12.3	14.6	9.3	17.6
Current marital status				
Not married	10.8	13.5	17.3	26.9
Currently married	89.2	86.5	82.7	73.1
Occupation				
Not working	35.5	1.1	27.7	0.2
Prof/cleric/sales/serv	8.4	43.3	10.3	15.7
Agriculture	42.1	66.7	43.3	58.0
Manual	14.0	20.7	18.7	26.1
Religion				
Catholic	51.5	47.8	42.2	46.2
Protestant	31.8	35.0	35.4	35.3
Adventist	12.1	12.6	14.6	12.8
Other	4.6	4.6	7.8	5.7
Type of place of residence				
Urban	16.2	15.9	15.7	21.3
Rural	83.8	84.1	84.3	78.7
Wealth				
Poorest	19.0	15.4	12.1	12.7
Poorer	19.1	17.8	29.0	16.6
Middle	20.7	21.5	24.4	25.1
Richer	21.3	21.8	16.7	19.8
Richest	20.0	23.5	17.8	25.8
Circumcision				
No	89.8	86.3	87.7	81.9
Yes	10.2	13.7	12.3	18.1

Trends in HIV Knowledge among Men

From 2005 to 2010, among all men who had sex in the year preceding the survey, basic knowledge of HIV appeared to decrease, while among men who had two or more partners in the year preceding the survey, it appeared to increase. However, neither change was statistically significant. The trends in the two other types of knowledge are similar among all men who had recent sex and among men with two or more partners. In both groups, knowledge of the protective effect of having only one uninfected faithful partner decreased, while knowledge of the protective effect of using condoms increased. The decrease in knowledge of the protective effect of one uninfected faithful partner and the increase in the knowledge of the protective effect of using condoms were only statistically significant among all men who had sex in the year preceding the survey. (Table 2)


Table 2. HIV-Related knowledge among study population

	Among those who had sex last year				Among those who had 2+ partners last year			
	<u>2005 (n=2,773)</u>		<u>2010 (n=3,772)</u>		<u>2005 (n=138)</u>		<u>2010 (n=262)</u>	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI
Basic knowledge of HIV	71.0	68.9-73.0	67.7	65.8-69.5	65.3	56.6-73.1	68.2	61.7-74.1
Knowledge of protective effect of one uninfected faithful partner	89.8	88.1-91.4	82.0	80.2-83.6	89.2	82.2-93.7	83.9	78.7-88.1
Knowledge of protective effect of using condoms	89.8	88.2-91.2	94.0	93.1-94.6	92.3	86.6-95.7	96.9	92.3-98.8

Risky Sexual Behavior by HIV Knowledge

In 2010, more men reported that they had two or more sexual partners compared with 2005 (7% versus 5%). However, among those who had two or more sexual partners, the percentage using a condom at last sex was higher in 2010 than 2005, at 93% versus 75%. The association is statistically significant (Figure 3).

Figure 3. Percent of men who had two types of risky sexual behavior



Note: Bars indicate 95% confidence intervals. Difference between years is statistically significant on both outcomes.

Table 3 shows that, among men with basic knowledge of HIV, knowledge of protective effect of one uninfected faithful partner, and knowledge of the protective effect of using condoms, a significantly higher proportion had two or more partners in 2010 compared with 2005.

In 2010, among all knowledge groups, a statistically significantly lower proportion of men with two or more partners reported not using a condom at the last sex than in 2005, meaning that risky sexual behavior declined between the two surveys (Table 3).

Table 3. Risky sexual behavior by knowledge of HIV, 2005 and 2010

	Men who had 2+ sexual partners last year				Men who did not use a condom at the last sex among those who had 2+ sexual partners			
	<u>2005 (n=2,773)</u>		<u>2010 (n=3,772)</u>		<u>2005 (n=138)</u>		<u>2010 (n=262)</u>	
	%	95% CI	%	95% CI	%	95% CI	%	95% CI
Basic knowledge of HIV	4.6	3.7-5.7	7.0	6.0-8.1	95.1	88.5-98.0	71.3	64.5-77.2
Knowledge of protective effect of one uninfected faithful partner	4.9	4.1-6.0	7.1	6.2-8.1	93.8	87.6-97.0	75.1	69.0-80.4
Knowledge of protective effect of using condoms	5.1	4.2-6.2	7.2	6.3-8.1	92.7	86.8-96.1	74.0	68.4-78.9

Association between HIV Knowledge and Risky Sexual Behavior

The study explored the association between HIV knowledge and risky sexual behavior using bivariate analysis and logistic regression. The model used is a pooled model of association between HIV knowledge and risky behavior.

Each type of HIV knowledge (basic knowledge, knowledge of the protective effect of one uninfected faithful partner, and knowledge of the protective effect of using condoms) was regressed separately with each risky sexual behavior (having two or more sexual partners; not using a condom at the last high risk sex) in the unadjusted models. Then, HIV knowledge variables and background variables were regressed together with a set of control factors in the adjusted models.

In unadjusted models, the only HIV knowledge variable associated with any risky sexual behavior outcome was knowledge of the protective effect of using condoms. Men who had this knowledge had higher odds of having two or more partners in the past year; interestingly, this knowledge was not significantly associated with condom use.

The year of the survey was not statistically significant in the adjusted model of having two or more partners, but it was statistically significant in the adjusted model of condom use, indicating that men with two or more partners in 2010 were significantly less likely to report not using a condom at last sex than men in 2005, even after adjusting for knowledge and characteristics.

The odds of having two or more sexual partners are greater among men in the poorer and middle wealth quintiles, while married men are less likely to have two or more sexual partners compared with unmarried men. However, even after adjusting for other factors, married men have 37 times the odds of not using a condom compared with unmarried men, when they have two or more sexual partners.

Professional occupations and manual workers have increased odds of having multiple sexual partners compared with men who are not working, but among those with more than two partners, men who work in professional occupations and manual labor have significantly lower odds of not using a condom. Rural-urban residence is unrelated to either type of risky behavior in the adjusted models.

Table 4. Pooled multivariate association between background factors, knowledge, and risky sexual behaviors 2005 and 2010

	Had 2+ sexual partners among those who had sex			Not using condom at last sex, among those with 2+ sexual partners				
	Unadjusted OR	CI	Adjusted OR	CI	Unadjusted OR	Adjusted OR	CI	
HIV Knowledge								
Basic knowledge of HIV	0.9	0.7-1.2	0.8	0.7-1.1	0.7	0.4-1.3	0.9	0.4-2.0
Knowledge of protective effect of one uninfected faithful partner	1.0	0.7-1.4	1.1	0.8-1.5	1.4	0.7-2.9	1.3	0.6-3.0
Knowledge of protective effect of using condoms	1.8 **	1.0-3.1	1.7	1.0-2.9	0.2	0.03-1.6	0.2	0.02-1.6
Year of survey								
2005 (ref.)								
2010	1.4 **	1.1-1.8	1.2	0.9-1.5	0.2 ***	0.1-0.4	0.2 ***	0.1-0.4
Age								
15-24 years (ref.)								
25-34 years	0.9	0.7-1.3	1.4 **	1.0-2.0	3.7 ***	2.0-7.1	1.2	0.5-2.9
35-44 years	0.7	0.5-1.0	1.4	1.0-2.1	6.5 ***	2.9-14.4	0.6	0.2-2.0
45-54 years	0.9	0.6-1.3	1.7 ***	1.1-2.7	10.7 ***	4.3-26.5	0.6	0.2-2.5
55-59 years	0.9	0.5-1.4	1.7 *	1.0-2.9	6.4 **	1.7-23.4	0.2 *	0.05-1.2
Occupation								
Not working (ref.)								
Prof/Cleric/Sales/Serv	2.3 ***	1.4-3.7	2.1 ***	1.3-3.5	0.05 ***	0.01-0.2	0.2	0.03-1.8
Agriculture	1.5 *	1.0-2.4	1.3	0.9-2.1	0.5	0.1-2.0	0.9	0.1-6.4
Manual	2.2 ***	1.4-3.5	1.8 **	1.2-2.9	0.01 **	0.03-0.5	0.5	0.1-3.7
Education level								
No education (ref.)								
Primary	1.5 **	1.1-2.0	1.5 **	1.1-2.0	0.5	0.2-1.4	0.5	0.1-2.6
Secondary	1.5 *	1.0-2.2	1.3	0.8-2.0	0.2 *	0.07-0.6	2.0	0.3-13.4

(Continued...)

Table 4. – Continued

	Had 2+ sexual partners among those who had sex		Not using condom at last sex, among those with 2+ sexual partners	
	Unadjusted OR	Adjusted OR	Unadjusted OR	Adjusted OR
Marital status				
Not married (ref.)				
Currently married	0.4 ***	0.4 ***	24.3 ***	37.1 ***
	0.3-0.5	0.3-0.5	13.2-44.8	13.3-103.2
Religion				
Catholic (ref.)				
Protestant	1.2	1.2	1.2	0.8
	0.9-1.5	0.9-1.5	0.7-2.1	0.4-1.9
Adventist	1.2	1.3	2.3	2.0
	0.9-1.7	0.9-1.8	0.9-6.2	0.4-9.4
Other	1.6 *	1.5 *	0.6	0.7
	1.0-2.5	1.0-2.4	0.2-1.3	0.2-2.9
Type of place of residence				
Urban (ref.)				
Rural	0.9	0.9	4.7 ***	1.5
	0.6-1.0	0.6 - 1.2	2.7-8.3	0.6-3.8
Wealth index				
Poorest (ref.)				
Poorer	1.6 *	1.6 **	4.6 *	0.9
	1.1-2.3	1.1 - 2.3	1.3-16.4	0.2-4.1
Middle	1.6 **	1.5 **	1.3	1.0
	1.1-2.4	1.0 - 2.2	0.5-3.3	0.3-3.9
Richer	1.2	1.1	1.5	0.7
	0.8-1.8	0.7 - 1.6	0.6-4.1	0.2-3.0
Richest	1.4	0.9	0.4	0.5
	1.0-2.1	0.6 - 1.4	0.2-0.8	0.1-2.1

Note: * = p<0.05; ** = p<0.01; *** = p<0.001

DISCUSSION AND CONCLUSIONS

This study explores HIV knowledge and risky sexual behavior. Using the 2005 and 2010 RDHS, we found a number of important results. First, among sexually active men, the findings show that knowledge of the protective effect of using condoms has significantly increased, comparing 2010 with 2005, and this is a very encouraging result. However, basic knowledge of HIV (knowing that a healthy-looking person can have the AIDS virus, knowing that HIV is not transmitted by mosquito bites or sharing food) has not significantly changed, and knowledge of the protective effect of having one uninfected faithful partner has decreased between 2005 and 2010, both among men who had sex in last year preceding the survey and men who had multiple partners. This result suggests the need for more HIV education particularly on the issues discussed above. Unfortunately, for the moment it is not possible to explain this decrease of HIV knowledge through HIV programmatic activities. In fact, it seems that there is no significant change in the implementation of HIV prevention activities since 2005 that could explain this decrease.

Our findings also show that in 2010 the percentage of men who used a condom at last sex significantly increased compared with 2005. In their study on HIV-related knowledge and behaviors in Rwanda, Rathavuth Hong and colleagues also found a considerable increase in condom use during last sex and during last non-spousal sex (Hong et al. 2013). This result is encouraging, and it may be the positive effect of HIV/AIDS prevention interventions that Rwanda has been implementing. For example, the observed increase in condom use could be linked to the significant increase of knowledge of the protective effect of using condoms (from 90% in 2005 to 94% in 2010).

In an unadjusted logistic regression model, sexually active men with knowledge of the protective effect of using condoms had 1.8 times higher odds of having two or more partners in the past year, a finding which may suggest some degree of reverse causation: men who engage in risky behaviors are more knowledgeable about the means of risk reduction. Otherwise the logistic regression analysis did not find any relationship between HIV knowledge and the two types of risky sexual behavior studied. Although a bit surprising, this finding corroborates the results of other studies (Essien et al. 2002; Williams et al. 2003). In their study on HIV/AIDS information and risky behavior in Botswana, Dinkelman et al. (2006) also concluded, “it may be

overly optimistic to hope for reductions in risky behavior through the channel of HIV-information provision alone.”

Therefore, this is a very important subject, particularly for policymakers who decide on strategies, interventions, and activities for HIV prevention. It is necessary not to focus only on AIDS knowledge as a means to reduce HIV transmission. Further studies could provide more information on other elements that probably influence safer sexual behavior and that therefore could be a focus for designing HIV prevention programs.

Our study has some limitations that should be considered when interpreting its results. Few men reported having two or more sexual partners (only 5% in 2005 and 7% in 2010). On one hand, this could be an under-reporting of sexual behavior. In fact, when men are interviewed in close quarters with other family members, they may not be willing to disclose their sexual behavior. On the other hand, the small sample size of men who had two or more sexual partners could be a reason for lack of significant association found between HIV knowledge and condom use. It is also possible to have recall bias in condom use among men who have not recently had sex.

In conclusion, the findings suggest that, while knowledge of the protective effect of condoms has increased, basic HIV knowledge has declined slightly among Rwandan men, and this should draw attention and lead to take appropriate measures. However, while setting up policies and strategies related to HIV prevention, policymakers should consider that HIV knowledge alone does not appear to be sufficient. Further research may help to identify other factors that are essential to consider.

REFERENCES

- Ahmed, S., T. Lutalo, M. Wawer, D. Serwadda, N.K. Sewankambo, F. Nalugoda, F. Makumbi, F. Wabwire-Mangen, N. Kiwanuka, G. Kigozi, M. Kiddugavu, and R. Gray. 2001. "HIV Incidence and Sexually Transmitted Disease Prevalence Associated with Condom Use: A Population Study in Rakai, Uganda." *AIDS* 15(16): 2171-9.
- Biraro, S., L.A. Shafer, I. Kleinschmidt, B. Wolff, A. Karabalinde, A. Nalwoga, J. Musinguzi, W. Kirungi, A. Optio, J. Whitworth, and H. Grosskurth. 2009. "Is Sexual Risk Taking Behaviour Changing in Rural South-West Uganda? Behaviour Trends in a Rural Population Cohort 1993-2006." *Sexually Transmitted Infections* 85(Suppl 1): i3-11.
- Booyesen, F. 2004. "HIV/AIDS, Poverty and Risky Sexual Behaviour in South Africa." *African Journal of AIDS Research* 3: 57-67.
- Cassell, M.M., D.T. Halperin, J.D. Shelton, and D. Stanton. 2006. "Risk Compensation: The Achilles' Heel of Innovations in HIV Prevention?" *BMJ* 332(7541): 605-7.
- Centers for Disease Control and Prevention (CDC). 2010. *Youth Risk Behavior Surveillance, United States, 2009*. Available at <http://www.cdc.gov/mmwr/preview/mmwrhtml/ss5905a1.htm>.
- de Walque, D., and R. Kline. 2009. *Comparing Condom Use with Different Types of Partners: Evidence from National HIV Surveys in Africa*. Washington D.C., USA: The World Bank.
- Dinkelman, T., J. Levinsohn, and R. Majelantle. 2006. *When Knowledge Is Not Enough: HIV/AIDS Information and Risky Behavior in Botswana*. National Bureau of Economic Research Working Paper 12418. Cambridge, Massachusetts, USA: NBER.
- Eaton, L., and S.C. Kalichman. 2009. "Behavioral Aspects of Male Circumcision for the Prevention of HIV Infection." *Curr HIV/AIDS Rep* 6(4): 187-93.
- Essien, E.J., O. Mgbere, E. Monjok, E. Ekong, S. Abughosh, and M.M. Holstad. 2002. "Predictors of Frequency of Condom Use and Attitudes among Sexually Active Female Military Personnel in Nigeria." *HIV AIDS (Auckl)* 2: 77-88.

- Gillespie, C. 1997. "Women's HIV Risk Reduction Efforts and Traditional Models of Health Behavior: A Review and Critique." *Women's Health* 3(1): 1-30.
- Gras, M.J., B.H. van Benthem, R.A. Coutinho, and A. van den Hoek. 2001. "Determinants of High Risk Sexual Behavior among Immigrant Groups in Amsterdam: Implications for Interventions." *Journal of Acquired Immune Deficiency Syndromes* 28(2): 166-72.
- Hong, R., J. de Dieu, J.U. Condo, M. Ribakare, and E. Murekatete. 2013. *Recent Trends in HIV-Related Knowledge and Behaviors in Rwanda, 2005-2010: Further Analysis of the Demographic and Health Surveys. DHS Further Analysis Reports No. 89*. Calverton, Maryland, USA: ICF International.
- ICF International (ICFI). 2012. *The DHS Program STATcompiler*. Available online at <http://www.statcompiler.com>
- Kibira, S.P., E. Nansubuga, N.M. Tumwesigye, L.M. Atuyambe, and F. Makumbi. 2014. "Differences in Risky Sexual Behaviors and HIV Prevalence of Circumcised and Uncircumcised Men in Uganda: Evidence from a 2011 Cross-Sectional National Survey." *Reprod Health Matters* 11(1): 25.
- Kimuna, S., and Y. Djamba. 2005. "Wealth and Extramarital Sex among Men in Zambia." *International Family Planning Perspectives* 31(2): 83-9.
- Kumari, A., and R.J. Nair. 2012. "Predictors of High Risk Sexual Behaviour among Men in India." *The Journal of Family Welfare* 58(2): 25-34.
- Lagarde, E., B. Auvert, J. Chege, T. Sukwa, J.R. Glynn, H.A. Weiss, E. Akam, M. Laourou, M. Carael, A. Buve, and Study Group on the Heterogeneity of HIV Epidemics in African Cities. 2001. "Condom Use and Its Association with HIV/Sexually Transmitted Diseases in Four Urban Communities of Sub-Saharan Africa." *AIDS* 15(Suppl 4): S71-8.
- MacIntyre, K., L. Brown, and S. Sosler. 2001. "It's Not What You Know, But Who You Knew: Examining the Relationship between Behavior Change and AIDS Mortality in Africa." *AIDS Education and Prevention* 13(2): 160-74.
- Magnani, R.J., A.M. Karim, L.A. Weiss, K.C. Bond, M. Lemba, and G.T. Morgan. 2002. "Reproductive Health Risk and Protective Factors among Youth in Lusaka, Zambia." *J Adolesc Health* 30(1): 76-86.

- McKusick, J., and T. Coates. 1985. "Reported Changes in the Sexual Behavior of Men at Risk for AIDS. San Francisco. The AIDS Behavioral Research Project." *Public health Rep* 100(6): 622-9.
- Ministry of Health (MOH) [Rwanda], and Rwanda Biomedical Center (RBC). 2009. *Rwanda National Strategic Plan on HIV and AIDS 2009-2012*. Kigali, Rwanda: MOH and RBC.
- Ministry of Health (MOH) [Rwanda], and Rwanda Biomedical Center (RBC). 2014. *Rwanda Global AIDS Response Progress Report (GARPR) 2014*. Kigali, Rwanda: MOH and RBC.
- National Institute of Statistics of Rwanda (NISR) [Rwanda], Ministry of Health (MOH) [Rwanda], and ICF International (ICFI). 2012. *Rwanda Demographic and Health Survey 2010*. Calverton, Maryland, USA: NISR, MOH, and ICF International.
- National Institute of Statistics of Rwanda (NISR) [Rwanda], Ministry of Health (MOH) [Rwanda], and ICF International (ICFI). 2006. *Rwanda Demographic and Health Survey 2005*. Calverton, Maryland, USA: NISR, MOH, and ICF International.
- Opio, A., V. Mishra, R. Hong, J. Musinguzi, W. Kirungi, A. Cross, J. Mermin, and R. Bunnell. 2008. "Trends in HIV-Related Behaviors and Knowledge in Uganda, 1989-2005: Evidence of a Shift toward More Risk-Taking Behaviors." *JAIDS* 49(3): 320-6.
- Pinkerton, S.D., and P.R. Abramson. 1997. "Effectiveness of Condoms in Preventing HIV Transmission." *Soc Sci Med* 44(9): 1303-12.
- President's Emergency Plan for AIDS Relief (PEPFAR). 2013. *Next Generation Indicators Reference Guide*. Available at <http://www.pepfar.gov/documents/organization/206097.pdf>
- Sahlu, T., E. Kassa, T. Agonafer, A. Tsegaye, T. Rinke de Wit, H. Gebremariam, R. Doorly, I. Spijkerman, H. Yeneneh, R.A. Coutinho, and A.L. Fontanet. 1999. "Sexual Behaviours, Perception of Risk of HIV Infection, and Factors Associated with Attending HIV Post-Test Counselling in Ethiopia." *AIDS* 13(10): 1263-72.
- Seidman, S.N., W.D. Mosher, and S.O. Aral. 1994. "Predictors of High Risk Behavior in Unmarried American Women: Adolescent Environment as Risk Factor." *J Adolescent Health* 15(2): 126-32.

- Li, S., H. Huang, Y. Cai, G. Xu, F. Huang, and X. Shen. 2009. "Characteristics and Determinants of Sexual Behavior among Adolescents of Migrant Workers in Shanghai (China)." *BMC Public Health* 9: 195.
- UNAIDS. 2008. *Report on the Global AIDS Epidemic*. Geneva, Switzerland: UNAIDS.
- Weller, S., and K. Davis. 2002. "Condom Effectiveness in Reducing Heterosexual HIV Transmission." *Cochrane Database Syst Rev* (1): CD003255.
- Williams, B.G., D. Taljaard, C.M. Campbell, E. Gouws, L. Ndhlovu, J. van Dam, M. Carael, and B. Auvert. 2003. "Changing Patterns of Knowledge, Reported Behaviour and Sexually Transmitted Infections in a South African Gold Mining Community." *AIDS* 17(14): 2099-107.