


Zambia



**Demographic and
Health Survey**

2001-2002

World Summit for Children Indicators, Zambia 2001-2002

Childhood mortality	Infant mortality rate	95 per 1,000
	Under-five mortality rate	168 per 1,000
	Maternal mortality rate	729 per 100,000
Childhood undernutrition	Percent stunted (children under 5 years)	46.8
	Percent wasted (children under 5 years)	5.0
	Percent underweight (children under 5 years)	28.1
Clean water supply	Percent of households with safe water supply ¹	51.3
Sanitary excreta disposal	Percent of households with flush toilets, pit toilet/latrine	70.2
Basic education	Proportion of children reaching grade 5 ²	87.7
	Net primary school attendance rate ²	67.1
	Proportion of children entering primary school ²	25.7
Family planning	Contraceptive prevalence rate (any method, currently married women)	34.2
	Contraceptive prevalence rate (any method, all women)	24.6
Antenatal care	Percent of women who received antenatal care from a health professional ³	93.4
Delivery care	Percent of births in the 5 years preceding the survey attended by a health professional	43.4
Low birth weight	Percent of births in the 5 years preceding the survey at low birth weight ⁴	10.7
Iodised salt intake	Percent of households that use iodised salt ⁵	79.8
Vitamin A supplements	Percent of children age 6-59 months who received a vitamin A dose in the 6 months preceding the survey	67.4
	Percent of women age 15-49 who received a vitamin A dose in the 2 months after delivery ³	27.5
Exclusive breastfeeding	Percent of youngest children under 6 months who are exclusively breastfed	40.1
Continued breastfeeding	Percent of children age 12-15 months still breastfeeding	96.8
	Percent of children age 20-23 months still breastfeeding	55.5
Timely complementary feeding	Percent of youngest children age 6-9 months receiving breast milk and complementary foods	87.4
Vaccinations	Percent of children age 12-23 months with BCG vaccination	94.0
	Percent of children age 12-23 months with at least 3 DPT vaccinations	80.0
	Percent of children age 12-23 months with at least 3 polio vaccinations	80.2
	Percent of children age 12-23 months with measles vaccination	84.4
	Percent of mothers who received at least 2 tetanus toxoid vaccinations during pregnancy ³	26.7
Oral rehydration therapy (ORT)	Percent of children age 0-59 months with diarrhoea in the 2 weeks preceding the survey who received oral rehydration salts (ORS)	53.2
Home management of diarrhoea	Percent of children age 0-59 months with diarrhoea in the 2 weeks preceding the interview who took more fluids than usual and continued eating somewhat less, the same or more food	28.3
Treatment of ARI		69.1
Treatment of illness	Percent of children age 0-59 months with acute respiratory infection (ARI) in the 2 weeks preceding the survey who were taken to a health provider	60.3
	Percent of children age 0-59 months with diarrhea, fever and/or ARI in the two weeks preceding the survey who were taken to a health provider	16.2
Children in especially difficult situations	Percent of children with at least one parent dead ²	15.0
	Percent of children who do not live with either biological parent ²	9.8
Use of impregnated bednets	Percentage of children 0-59 months who slept under an impregnated bednet on the previous night ⁶	51.9
Malaria treatment	Percent of children age 0-59 months with a fever in the 2 weeks preceding the survey who were treated with an anti-malarial drug	66.3
		43.3
HIV/AIDS	Percent of women age 15-49 who correctly stated 2 ways of avoiding HIV infection ⁷	55.4
	Percent of women age 15-49 who correctly identified 2 misconceptions about HIV/AIDS ⁸	64.4
	Percent of women age 15-49 who believe that AIDS can be transmitted from mother to child during pregnancy, delivery and breastfeeding	9.4
	Percent of women age 15-49 who know of a place to get tested for the AIDS virus	
	Percent of women age 15-49 who have been tested for the AIDS virus	

¹Piped water or protected well water

²Based on de jure children

³For the last live birth in the five years preceding the survey

⁴For children without a reported birth weight, the proportion with low birth weight is assumed to be the same as the proportion with low birth weight in each birth size category among children who have a reported birth weight.

⁵15 parts per million or more

⁶Refers to ever-impregnated bednets

⁷Having sex with only one partner who has no other partners and using a condom every time they have sex

⁸They say that AIDS cannot be transmitted through mosquito bites and that a healthy-looking person can have the AIDS virus.

Zambia

Demographic and Health Survey

2001-2002

Central Statistical Office
Lusaka, Zambia

Central Board of Health
Lusaka, Zambia

ORC Macro
Calverton, Maryland, USA

February 2003



Central
Statistical Office



Central Board
of Health



ORC Macro

This report summarises the findings of the 2001-2002 Zambia Demographic and Health Survey (ZDHS) carried out by the Central Statistical Office in partnership with the Central Board of Health. ORC Macro provided financial and technical assistance for the survey through the USAID-funded MEASURE *DHS+* programme, which is designed to assist developing countries to collect data on fertility, family planning, and maternal and child health. Additional funding for the ZDHS was received from the Government of Japan through a trust fund managed by the United Nations Development Programme (UNDP) and through bilateral agreements between the Government of the Republic of Zambia and the United Nations Population Fund (UNFPA) and the Danish International Development Agency (DANIDA). The opinions expressed are those of the authors and do not necessarily reflect the views of USAID, UNDP, UNFPA or DANIDA.

Additional information about the ZDHS may be obtained from the Central Statistical Office, P.O. Box 31908, Lusaka, Tel: 260-1-251377/85; fax: 260-1253468. Additional information about the MEASURE *DHS+* project may be obtained by contacting: MEASURE *DHS+*, ORC Macro, 11785 Beltsville Drive, Suite 300, Calverton, MD 20705 (telephone 301-572-0200; fax 301-572-0999; e-mail: reports@macroint.com; internet: www.measuredhs.com).

Suggested citation:

Central Statistical Office [Zambia], Central Board of Health [Zambia], and ORC Macro. 2003. *Zambia Demographic and Health Survey 2001-2002*. Calverton, Maryland, USA: Central Statistical Office, Central Board of Health, and ORC

Macro.

CONTENTS

	Page
Tables and Figures	xi
Preface	xix
Summary of Findings	xxi
Map of Zambia.....	xxvi

CHAPTER 1 INTRODUCTION

Kumbutso Dzekedzeke and Chanda Mulenga

1.1	History, Geography, and Economy	1
1.2	Population	2
1.3	The Population Policy and National Population and Development Programme.....	3
1.4	Health Priorities and Programmes.....	4
1.6	Objectives and Organisation of the Survey.....	5
1.7	Sample Design.....	6
1.8	Questionnaires	6
1.9	Syphilis and HIV Testing	7
1.10	Pretest Activities, Training, and Fieldwork	8
1.11	Data Processing	10
1.12	Response Rates.....	10

CHAPTER 2 CHARACTERISTICS OF HOUSEHOLDS AND RESPONDENTS

Patrick Mumba Chewe

2.1	Population by Age and Sex	11
2.2	Household Composition	13
2.3	Children's Living Arrangements and Parental Survival	14
2.4	Educational Level of Household Population	15
2.5	Housing Characteristics and Household Possessions.....	21
2.6	Household Durable Goods	22
2.7	Access to Health Care.....	23

CHAPTER 3 CHARACTERISTICS OF RESPONDENTS AND WOMEN'S CHARACTERISTICS

Sheila S. Mudenda

3.1	Characteristics of Survey Respondents	25
3.2	Educational Attainment.....	25
3.3	Literacy.....	29
3.4	Access to Mass Media	31
3.5	Men's Employment.....	33

	Page
3.6	Women’s Employment36
3.6.1	Employment Status36
3.6.2	Occupation37
3.6.3	Earnings, Type of Employer, and Continuity of Employment38
3.6.4	Control over Women’s Earnings and Contribution to Household Expenditures.....39
3.7	Women’s Empowerment41
3.7.1	Women’s Participation in Decision-making.....41
3.7.2	Attitudes toward Wife Beating46
3.7.3	Attitudes toward Refusing Sex.....49
3.8	Life-style Measures52
3.8.1	Use of Tobacco52
CHAPTER 4	FERTILITY
	<i>Richard Banda and Margaret T. Mwanamwenge</i>
4.1	Fertility Levels and Trends.....55
4.1.1	Fertility Levels.....55
4.1.2	Differentials in Current and Completed Fertility57
4.1.3	Trends in Fertility.....59
4.2	Children Ever Born and Children Surviving62
4.3	Birth Intervals64
4.4	Age at First Birth65
4.5	Median Age at First Birth by Background Characteristics66
4.6	Teenage Fertility67
CHAPTER 5	FERTILITY REGULATION
	<i>Mika Bwembya and Margaret T. Mwanamwenge</i>
5.1	Knowledge of Contraceptive Methods69
5.2	Ever Use of Contraception71
5.3	Current Use of Contraceptive Methods.....74
5.4	Current Use of Contraception by Background Characteristics.....75
5.5	Trends in the Use of Family Planning77
5.6	Number of Children at First Use of Contraception78
5.7	Use of Social Marketing Brand Pills.....79
5.8	Knowledge of Fertile Period.....79
5.9	Source of Supply80

	Page
5.10	Informed Choice81
5.11	Future Use of Contraception.....82
5.12	Reasons for Not Intending to Use Contraception83
5.13	Preferred Method of Contraception for Future Use84
5.14	Exposure to Family Planning Messages.....85
5.15	Exposure to Specific Radio and Television Shows on Family Planning and HIV86
5.16	Contact of Non-users with Family Planning Providers.....88
5.17	Discussion about Family Planning with Husband89
5.18	Attitudes of Couples toward Family Planning90
CHAPTER 6	OTHER DETERMINANTS OF FERTILITY
	<i>Anne R. Cross</i>
6.1	Current Marital Status93
6.2	Polygyny94
6.3	Age at First Marriage95
6.4	Age at First Sexual Intercourse97
6.5	Recent Sexual Activity.....99
6.6	Postpartum Amenorrhoea, Abstinence and Insusceptibility 102
6.7	Menopause 103
CHAPTER 7	FERTILITY PREFERENCES AND UNMET NEED FOR FAMILY PLANNING
	<i>Margaret T. Mwanamwenge and Richard Banda</i>
7.1	Desire for More Children..... 105
7.2	Desire to Limit Childbearing by Background Characteristics 107
7.3	Unmet Need for Family Planning..... 109
7.4	Ideal Family Size..... 111
7.5	Wanted and Unwanted Fertility..... 114
CHAPTER 8	INFANT AND CHILD MORTALITY
	<i>Kumbutso Dzikedzeke</i>
8.1	Introduction 117
8.2	Levels and Trends in Infant and Child Mortality 118
8.3	Early Childhood Mortality by Socioeconomic Characteristics..... 119
8.4	Early Childhood Mortality Rates by Demographic Characteristics 121
8.5	Women’s Status and Early Childhood Mortality 122
8.6	High-Risk Fertility Behaviour..... 123

CHAPTER 9 MATERNAL AND CHILD HEALTH*Miriam Chipimo, Elizabeth Mulamfu, Martha Mulenga, and Arlinda Zhuzhuni*

9.1	Maternity Care	127
9.1.1	Antenatal Care.....	127
9.1.2	Delivery and Postnatal Care.....	132
9.1.3	Trends in Maternity Care Indicators	137
9.2	Visits by Health Workers and Presence of a Neighbourhood Health Committee ...	138
9.3	Women’s Perceptions of Problems in Obtaining Health Care.....	141
9.4	Child Immunisation	143
9.5	Acute Respiratory Infections	147
9.6	Diarrhoeal Diseases	149

CHAPTER 10 MALARIA*Miriam Chipimo and Richard Banda*

10.1	Introduction	155
10.1.1	Ownership of Mosquito Nets.....	155
10.1.1	Use of Bednets by Children and Pregnant Women	156
10.2	Malaria Diagnosis, Case Management and Treatment	157
10.2.1	Malaria Prophylaxis during Pregnancy.....	157
10.2.2	Prevalence and Management of Childhood Malaria.....	158

CHAPTER 11 INFANT FEEDING, NUTRITIONAL PRACTICES, AND NUTRITIONAL STATUS AMONG YOUNG CHILDREN AND WOMEN*Bornwell Sikateyo, Agnes Mugala Aongola, and Arlinda Zhuzhuni*

11.1	Breastfeeding.....	163
11.1.1	Initiation of Breastfeeding	163
11.1.2	Age Pattern of Breastfeeding.....	165
11.2	Complementary Feeding.....	169
11.2.1	Types of Complementary Foods.....	169
11.2.2	Frequency of Foods Consumed by Children	171
11.3	Micronutrients	172
11.3.1	Salt and Sugar Fortification	172

	Page
11.3.2 Micronutrient Intake among Children	175
11.3.3 Micronutrient Supplementation for Women	177
11.4 Household Food Security.....	178
11.5 Nutritional Status of Children under Age Five.....	179
11.5.1 Measures of Nutritional Status in Childhood	179
11.5.2 Trends in Children’s Nutritional Status	183
11.6 Nutritional Status of Women	183

CHAPTER 12 VIOLENCE AGAINST WOMEN

Mary Kazunga and Patrick Mumba Chew

12.1 Introduction	185
12.2 Characteristics of Respondents.....	186
12.3 Experience of Beatings or Physical Mistreatment.....	186
12.4 Perpetrators of Physical Violence	189
12.5 Frequency of Beatings.....	189
12.6 Sexual Violence by a Man.....	190
12.7 Perpetrators and Frequency of Sexual Violence in the Past 12 Months.....	192
12.8 Sexual Intercourse Forced by a Third Party	194

CHAPTER 13 HIV/AIDS/STI-RELATED KNOWLEDGE AND BEHAVIOUR

Margaret T. Mwanamwenge

13.1 Introduction	195
13.2 Knowledge about AIDS and Ways to Avoid It	195
13.2.1 Knowledge of Ways to Avoid AIDS	195
13.2.2 Knowledge of Programmatically Important Ways to Avoid HIV/AIDS	197
13.3 Knowledge of HIV/AIDS Transmission and its Effects.....	199
13.4 Perception of HIV/AIDS.....	200
13.4.1 Discussion of HIV/AIDS with Partners	200
13.4.2 Stigma Associated with HIV/AIDS	201
13.4.3 Discussions of Condoms in the Media.....	203
13.5 Testing for HIV	205
13.6 Men’s Attitudes Toward Condoms.....	208
13.7 Absence from Home and Use of Alcohol	210
13.7.1 Absence from Home.....	210
13.7.2 Alcohol Consumption	211
13.8 Knowledge of Symptoms of Sexually Transmitted Infections.....	213

	Page
13.9	Reports of Recent Sexually Transmitted Infections 216
13.10	Treatment Seeking and Protection of a Partner from STIs 217
13.11	Sexual Behaviour 219
13.11.1	Number of Sexual Partners 219
13.11.2	Knowledge of Condom Sources and Condom Use for Disease Prevention 221
13.11.3	Payment for Sexual Relations 224
CHAPTER 14 PREVALENCE OF HIV AND SYPHILIS	
<i>Kumbutso Dzekedzeke</i>	
14.1	Coverage of HIV and Syphilis Testing 225
14.1.1	Coverage of Syphilis Testing 225
14.1.2	Coverage of HIV Testing 226
14.2	Syphilis Prevalence Rates 229
14.3	HIV Prevalence Rates 234
14.4	Comparison of ANC Surveillance Estimates and the ZDHS Rates 238
CHAPTER 15 ADULT MORTALITY	
<i>Ann A. Way</i>	
15.1	Adult Mortality Data 241
15.1.1	Data Collection Procedures 241
15.1.2	Data Quality Assessment 241
15.2	Direct Estimates of Adult Mortality 242
15.2.1	Levels and Trends in Adult Mortality 242
15.2.2	Differentials in Adult Mortality Levels 244
15.3	Estimates of Maternal Mortality 245
REFERENCES 247	
APPENDIX A SAMPLE DESIGN 251	
A.1	Introduction 251
A.2	Survey Objectives 251
A.3	Sample Domains 251
A.4	Sample Frame 251
A.5	Stratification 252
A.6	Sample Allocation 252

	Page
A.7 Sample Selection	254
A.8 Response Rates.....	254
APPENDIX B SAMPLING ERRORS	259
APPENDIX C DATA QUALITY TABLES	275
APPENDIX D SURVEY PERSONNEL.....	281
APPENDIX E QUESTIONNAIRES	285

TABLES AND FIGURES

		Page
CHAPTER 1	INTRODUCTION	
Table 1.1	Demographic characteristics	2
Table 1.2	Results of the household and individual interviews	10
CHAPTER 2	CHARACTERISTICS OF HOUSEHOLDS AND RESPONDENTS	
Table 2.1	Household population by age, sex, and residence	11
Table 2.2	Household composition.....	13
Table 2.3	Children’s living arrangements and orphanhood	14
Table 2.4	Educational attainment of household population	16
Table 2.5	School attendance ratios.....	18
Table 2.6	Grade repetition and dropout rates.....	20
Table 2.7	Household characteristics	21
Table 2.8	Household durable goods.....	23
Table 2.9	Inability to pay for medical care.....	24
Figure 2.1	Population pyramid	12
Figure 2.2	Distribution of de facto household population by five-year age groups and sex.....	13
Figure 2.3	Percent distribution of de facto household population with no education by age and sex	17
Figure 2.4	Net and gross attendance ratios by sex.....	19
Figure 2.5	Trends in percentage of households with electricity by residence, Zambia 1992-2002.....	22
Figure 2.6	Trends in percentage of households owning specific durable goods Zambia 1992-2002.....	23
CHAPTER 3	CHARACTERISTICS OF RESPONDENTS AND WOMEN’S CHARACTERISTICS	
Table 3.1	Background characteristics of respondents.....	26
Table 3.2	Educational attainment by background characteristics.....	28
Table 3.3	Literacy.....	30
Table 3.4	Exposure to mass media	32
Table 3.5	Employment status: men	33
Table 3.6	Occupation: men	35
Table 3.7	Type of earnings: men	36
Table 3.8	Employment status: women.....	37
Table 3.9	Occupation: women.....	38
Table 3.10	Employment characteristics: women	39

	Page
Table 3.11	Decision on use of earnings and contribution of earnings to household expenditures..... 40
Table 3.12	Women’s control over earnings 41
Table 3.13	Women’s participation in decisionmaking 42
Table 3.14.1	Women’s participation in decision-making by background characteristics: women 43
Table 3.14.2	Women’s participation in decision-making by background characteristics: men 45
Table 3.15.1	Women’s attitudes toward wife beating..... 47
Table 3.15.2	Men’s attitudes toward wife beating 48
Table 3.16.1	Women’s attitudes toward wife refusing sex with husband..... 50
Table 3.16.2	Men’s attitudes toward wife refusing sex with husband 51
Table 3.17.1	Smoking of tobacco: women 53
Table 3.17.2	Smoking of tobacco: men 53
Table 3.18	Smoking of tobacco by background characteristics..... 54
Figure 3.1	Youth literacy 31
Figure 3.2	Distribution of unemployed men by current activity 34
Figure 3.3	Percentage of women and men who agree with specific reasons that a wife is justified in refusing sex with her husband 52

CHAPTER 4 FERTILITY

Table 4.1	Current fertility 56
Table 4.2	Fertility by background characteristics..... 58
Table 4.3	Trends in age-specific and total fertility rates, various sources..... 59
Table 4.4	Trends in fertility by background characteristics 61
Table 4.5	Trends in age-specific fertility rates 62
Table 4.6	Children ever born and living..... 62
Table 4.7	Birth intervals..... 64
Table 4.8	Age at first birth 65
Table 4.9	Median age at first birth by background characteristics..... 66
Table 4.10	Teenage pregnancy and motherhood 67
Figure 4.1	Total fertility rates, selected sub-Saharan countries , 1998-2001 57
Figure 4.2	Total fertility rate by education level 58
Figure 4.3	Total fertility rate by province 59
Figure 4.4	Trends in age-specific fertility rates 1980 census, 1992 ZDHS, and 2001-2002 ZDHS..... 60
Figure 4.5	Trends in the total fertility rates by urban-rural residence, 1992 ZHDS, 1996 ZDHS, and 2001-2002 ZDHS 61
Figure 4.6	Percent distribution of all men and currently married men by number of children ever born, according to age group..... 63
Figure 4.7	Percentage of teenagers who are mothers or pregnant with their first child..... 68

CHAPTER 5 FERTILITY REGULATION

Table 5.1.1	Knowledge of contraceptive methods: women 70
Table 5.1.2	Knowledge of contraceptive methods: men 71

	Page
Table 5.2	Ever use of contraception: women..... 72
Table 5.3	Ever use of contraception: men 73
Table 5.4	Current use of contraception 75
Table 5.5	Current use of contraception by background characteristics..... 76
Table 5.6	Trends in the use of family planning..... 77
Table 5.7	Number of children at first use of contraception 79
Table 5.8	Use of Safeplan 79
Table 5.9	Knowledge of fertile period..... 80
Table 5.10	Source of contraception..... 81
Table 5.11	Informed choice 81
Table 5.12	Future use of contraception 82
Table 5.13	Reasons for not intending to use contraception..... 83
Table 5.14	Preferred method of contraception for future use 84
Table 5.15	Exposure to family planning messages..... 85
Table 5.16.1	Exposure to specific radio and television shows on family planning and HIV: Women..... 87
Table 5.16.2	Exposure to specific radio and television shows on family planning and HIV: Men 88
Table 5.17	Contact of non-users with family planning providers..... 89
Table 5.18	Discussion of family planning with husband..... 90
Table 5.19	Attitudes of couples toward family planning..... 91
Figure 5.1	Current use of any contraceptive method among currently married women age 15-49, by background characteristics 77
Figure 5.2	Trends in current use of specific contraceptive methods among currently married women age 15-49, Zambia 1992-2002..... 78

CHAPTER 6

OTHER DETERMINANTS OF FERTILITY

Table 6.1	Current marital status..... 93
Table 6.2	Polygyny..... 95
Table 6.3	Age at first marriage 96
Table 6.4	Median age at first marriage..... 97
Table 6.5	Age at first sexual intercourse..... 98
Table 6.6	Median age at first sexual intercourse 99
Table 6.7.1	Recent sexual activity: women..... 100
Table 6.7.2	Recent sexual activity: men 101
Table 6.8	Postpartum amenorrhoea, abstinence and insusceptibility 102
Table 6.9	Median duration of postpartum insusceptibility by background characteristics 103
Table 6.10	Menopause 103

CHAPTER 7

FERTILITY PREFERENCES AND UNMET NEED FOR FAMILY PLANNING

Table 7.1	Fertility preferences by number of living children..... 106
Table 7.2	Desire to limit childbearing..... 108
Table 7.3	Need for family planning 110
Table 7.4	Ideal and actual number of children 112
Table 7.5	Mean ideal number of children by background characteristics 113
Table 7.6	Fertility planning status 114

	Page
Table 7.7	Wanted fertility rates 115
Figure 7.1	Fertility preferences of currently married women 15-49..... 105
Figure 7.2	Percentage of married women and men who want no more children by number of living children 107
Figure 7.3	Percentage of women with three living children who do not want additional children, by background characteristics 108
Figure 7.4	Unmet need for spacing and limiting by residence and region 111
CHAPTER 8	INFANT AND CHILD MORTALITY
Table 8.1	Early childhood mortality rates..... 118
Table 8.2	Trends in early childhood mortality rates 119
Table 8.3	Early childhood mortality by background characteristics..... 120
Table 8.4	Trends in early childhood mortality rates by residence and education..... 121
Table 8.5	Early childhood mortality by demographic characteristics..... 122
Table 8.6	Early childhood mortality by women's status indicators 123
Table 8.7	High-risk fertility behaviour..... 124
CHAPTER 9	MATERNAL AND CHILD HEALTH
Table 9.1	Antenatal care 128
Table 9.2	Number of antenatal care visits and timing of first visit 129
Table 9.3	Antenatal care content..... 130
Table 9.4	Tetanus toxoid injections 131
Table 9.5	Place of delivery 133
Table 9.6	Assistance during delivery 134
Table 9.7	Delivery characteristics 135
Table 9.8	Postnatal care 137
Table 9.9	Visits by health workers 139
Table 9.10.1	Neighbourhood Health Committee (NHC): women..... 140
Table 9.10.2	Neighbourhood Health Committee (NHC): men 141
Table 9.11	Problems in accessing health care 142
Table 9.12	Vaccinations by source of information 143
Table 9.13	Vaccinations by background characteristics..... 145
Table 9.14	Prevalence of injections 147
Table 9.15	Prevalence and treatment of symptoms of acute respiratory infection (ARI)..... 148
Table 9.16	Prevalence of diarrhoea 149
Table 9.17	Knowledge of ORS packets..... 150
Table 9.18	Diarrhoea treatment 151
Table 9.19	Feeding practices during diarrhoea 152
Table 9.20	Household knowledge and usage of Clorin 153
Figure 9.1	Trends in maternity care indicators, Zambia 1992-2002 138
Figure 9.2	Trends in vaccination coverage among children 12-23 months 146
CHAPTER 10	MALARIA
Table 10.1	Ownership of mosquito nets..... 155
Table 10.2	Use of mosquito nets by children..... 156

	Page
Table 10.3	Use of mosquito nets by women..... 157
Table 10.4	Use of antimalarial drugs during pregnancy..... 158
Table 10.5	Prevalence and treatment of fever/convulsions in children 159
Table 10.6	Drugs taken for fever 160
Table 10.7	Treatment in children with fever/convulsions 161
CHAPTER 11	INFANT FEEDING, NUTRITIONAL PRACTICES, AND NUTRITIONAL STATUS AMONG YOUNG CHILDREN AND WOMEN
Table 11.1	Initial breastfeeding 164
Table 11.2	Breastfeeding status by child's age 166
Table 11.3	Median duration of breastfeeding 168
Table 11.4	Foods consumed by children in the day or night preceding the interview 170
Table 11.5	Frequency of foods consumed by children in the day or night preceding the interview 171
Table 11.6	Iodisation of household salt 173
Table 11.7	Presence of sugar in household..... 174
Table 11.8	Household consumption of sugar..... 174
Table 11.9	Micronutrient intake among children 176
Table 11.10	Micronutrient intake among mothers 178
Table 11.11	Household food security 179
Table 11.12	Nutritional status of children 181
Table 11.13	Nutritional status of women by background characteristics..... 184
Figure 11.1	Median duration of breastfeeding by background characteristics..... 167
Figure 11.2	Nutritional status of children under five years by province 182
CHAPTER 12	VIOLENCE AGAINST WOMEN
Table 12.1	Background characteristics of respondents who were asked questions about domestic violence 186
Table 12.2	Experience of beatings or physical mistreatment 187
Table 12.3	Perpetrators of physical violence..... 188
Table 12.4	Frequency of beatings..... 190
Table 12.5	Experience of sexual violence 191
Table 12.6	Perpetrators of sexual violence..... 193
Figure 12.1	Experience of sexual violence among ever-married women 192
Figure 12.2	Frequency of sexual violence in the past year, among women who report having ever experienced sexual violence 193
CHAPTER 13	AIDS/HIV/STI-RELATED KNOWLEDGE AND BEHAVIOUR
Table 13.1	Knowledge of HIV/AIDS 196
Table 13.2	Knowledge of ways to avoid HIV/AIDS 197
Table 13.3	Knowledge of programmatically important ways to avoid HIV/AIDS..... 198
Table 13.4	Knowledge of HIV/AIDS-related issues..... 200

	Page
Table 13.5	Discussion of HIV/AIDS with partner 201
Table 13.6	Social aspects of HIV/AIDS 202
Table 13.7	Discussion of condoms in the media..... 204
Table 13.8	Knowledge of “Trendsetters” newspaper..... 205
Table 13.9.1	Testing for the AIDS virus: women..... 206
Table 13.9.2	Testing for the AIDS virus: men..... 208
Table 13.10	Men's attitudes toward condoms..... 209
Table 13.11	Frequency and length of travel away from home 210
Table 13.12.1	Alcohol consumption: women..... 211
Table 13.12.2	Alcohol consumption: men..... 212
Table 13.13	Alcohol consumption by background characteristics 213
Table 13.14.1	Knowledge of symptoms of STDs: women 214
Table 13.14.2	Knowledge of symptoms of STDs: men..... 215
Table 13.15	Self-reporting of sexually transmitted infections and STI symptoms..... 216
Table 13.16	Source of treatment of STIs..... 217
Table 13.17	Efforts to protect partners from infection among women and men with STI 218
Table 13.18	Number of extramarital sexual partners: married women and men 219
Table 13.19	Number of sexual partners: unmarried women and men 220
Table 13.20	Knowledge of source for male condoms, and access to condoms..... 222
Table 13.21	Use of condoms by type of partner 223
Table 13.22	Commercial sexual activity and condom use during last commercial sex..... 224
Figure 13.1	Unmet need for HIV testing by education level..... 207
Figure 13.2	Percentage of men and women with an STI, discharge, genital sore, or ulcer..... 217
Figure 13.3	Percentage of unmarried women and men with at least two partners..... 221

CHAPTER 14 PREVALENCE OF HIV AND SYPHILIS

Table 14.1	Coverage for syphilis testing by urban-rural residence (unweighted) 225
Table 14.2	Coverage of syphilis testing by province (unweighted)..... 226
Table 14.3	Coverage for HIV testing by urban-rural residence (unweighted) 227
Table 14.4	Coverage of HIV testing by province (unweighted)..... 228
Table 14.5	Percentage of respondents tested for HIV 229
Table 14.6	Syphilis testing by age 231
Table 14.7	Syphilis prevalence by background characteristics..... 232
Table 14.8	Syphilis prevalence by age, sex, and urban-rural residence 233
Table 14.9	Syphilis prevalence by reproductive and sexual activity characteristics 234
Table 14.10	Syphilis prevention and treatment experience..... 235
Table 14.11	HIV prevalence by background characteristics 236
Table 14.12	HIV prevalence by age, sex, and urban-rural residence 238
Figure 14.1	Percentage of respondents 15-49 tested for HIV in the 2001-2002 ZDHS, by age group..... 230
Figure 14.2	HIV prevalence by sex and age 237
Figure 14.3	HIV prevalence among women by age and residence 239
Figure 14.4	HIV prevalence among men by age and residence..... 239

		Page
CHAPTER 15	ADULT MORTALITY	
Table 15.1	Completeness of reporting on siblings	242
Table 15.2	Adult mortality rates	243
Table 15.3	Adult mortality rates by residence and province	244
Table 15.4	Direct estimates of maternal mortality.....	245
Figure 15.1	Trends in Age-specific Mortality among Adults 15-49.....	244
APPENDIX A	SAMPLE DESIGN	
Table A.1	Proportional allocation and squared root allocation of 320 clusters by province	252
Table A.2	Expected number of selected households to reach the target of completed interviews by province	253
Table A.3	Final allocation of women 15-49 with completed interviews and clusters in each province by urban and rural residence	253
Table A.4	Sample implementation: women's sample	256
Table A.5	Sample implementation: women's sample	257
APPENDIX B	ESTIMATES OF SAMPLING ERRORS	
Table B.1	List of selected variables for sampling errors.....	261
Table B.2	Sampling errors for selected variables, total sample, Zambia	262
Table B.3	Sampling errors for selected variables, urban sample.....	263
Table B.4	Sampling errors for selected variables, rural sample.....	264
Table B.5	Sampling errors for selected variables, Central province	265
Table B.6	Sampling errors for selected variables, Copperbelt province	266
Table B.7	Sampling errors for selected variables, Eastern province	267
Table B.8	Sampling errors for selected variables, Luapula province	268
Table B.9	Sampling errors for selected variables, Lusaka province.....	269
Table B.10	Sampling errors for selected variables, Northern province	270
Table B.11	Sampling errors for selected variables, North-Western province	271
Table B.12	Sampling errors for selected variables, Southern province	272
Table B.13	Sampling errors for selected variables, Western province	273
APPENDIX C	DATA QUALITY TABLES	
Table C.1	Household age distribution	275
Table C.2.1	Age distribution of eligible and interviewed women.....	276
Table C.2.2	Age distribution of eligible and interviewed men	276
Table C.3	Completeness of reporting.....	277
Table C.4	Births by calendar years	277
Table C.5	Reporting of age at death in days	278
Table C.6	Reporting of age at death in months	279

PREFACE

The Zambia Demographic and Health Surveys is an important part of the surveillance system providing indicators for the strategic management and monitoring of the health sector in Zambia. Indicators from the surveys show the current status and trends in important areas of health care delivery such as environmental health, reproductive health, maternal health, child health, sexually transmitted infections and HIV/AIDS. The surveys are also a major source of information about health seeking behaviour regarding individuals and communities. Fertility and mortality indicators, which gauge the overall health status of the population are also provided by the surveys.

Three Demographic and Health Surveys have been carried out in Zambia, the first in 1992, the second in 1996, and the most recent in 2001-2002. Institutional collaboration has been the hallmark of the implementation of these surveys. As in the previous surveys, the Central Board of Health mobilised resources for the 2001-2002 ZDHS while the Central Statistical Office played the key role in the implementation of the survey. Other participating institutions were the Tropical Diseases Research Centre, which was in charge of syphilis and HIV testing; the University Teaching Hospital, which made important contributions towards the development of the syphilis and HIV testing protocol; the National Food and Nutrition Commission, which was responsible for the nutrition components; and the University of Zambia Demography Division, which assisted with the training of field staff.

Key people in the implementation of the 2001-2002 ZDHS were Mr David Diangamo, Director of Census and Statistics; Dr Musonda Rosemary Sunkutu, Director of Public Health and Research in the Central Board of Health; Mr Bornwell Sikateyo, Manager for Health Management Information Systems in the Central Board of Health; Mr Kumbutso Dzekedzeke, Survey Co-ordinator from the Central Statistical Office; Mr Patrick Mumba Chewe, Assistant Survey Co-ordinator from the Central Statistical Office; Dr Rosemary Musonda, Deputy Director at the Tropical Diseases Research Centre; and Ms Arlinda Zhuzhuni, Project Manager from ORC Macro.

A number of donors contributed towards the survey costs. The Government of Japan provided vehicles for the survey teams in addition to providing financial support for other components of the survey. Part of the support from the Government of Japan was channelled through the United Nations Development Programme. Other donors who contributed were the U.S. Agency for International Development, which provided financial and technical support through ORC Macro, the United Nations Population Fund, and the Danish Agency for International Development.

We owe an immense gratitude to the Field Coordinators, Interviewers, Nurse/Nurse Counsellors, Laboratory Technicians, Supervisors, Field Editors, Provincial Statistical Officers, and Drivers for their hard work and dedication. We have printed a list of names in Appendix D as a sign of our appreciation for their help and kindness. We are also grateful to all the respondents for their patience and generosity with their time.

It will only be worth the effort to have compiled all the indicators in this report if stakeholders in the health sector use them to improve health care delivery and efforts to prevent the spread of HIV and other infectious diseases in Zambia.



Dr. Ben U. Chirwa
Director General
Central Board of Health

SUMMARY OF FINDINGS

The 2001-2002 Zambia Demographic and Health Survey (ZDHS) was carried out by the Central Statistical Office and the Central Board of Health. It is a nationally representative sample of 7,658 women age 15-49 and 2,145 men age 15-59. The principal objective of the survey was to provide data to policymakers and planners on the population and health situation in Zambia. Most of the information collected in the 2001-2002 ZDHS represents updated estimates of basic demographic and health indicators covered in the 1992 ZDHS and 1996 ZDHS surveys. Specifically, the 2001-2002 ZDHS collected detailed information on fertility and family planning, child mortality and maternal mortality, maternal and child health and nutritional status, and knowledge, awareness and behaviour regarding HIV/AIDS and other sexually transmitted infections. New features of the 2001-2002 ZDHS include the collection of information on violence against women and testing of individuals for HIV and syphilis.

FERTILITY

Survey results indicate that the total fertility rate (TFR) in Zambia is 5.9. This means that at current fertility levels, the average Zambian woman who is at the beginning of her childbearing years will give birth to 5.9 children by the end of her reproductive period. Results from the 2001-2002 ZDHS show that fertility is highest in Northern province (6.9 births per woman) and lowest in Lusaka (4.3 births per woman). Education has a marked effect on fertility, with uneducated women having three and one-half more births than women with at least some secondary education (7.4 births versus 3.9 births).

Zambia's fertility continues to decline although comparatively slowly. The TFR has declined from a level of 6.5 births per woman in 1992 and 6.1 births per woman in 1996 to the current level of 5.9. Despite the decline, fertility in Zambia remains one of the highest in sub-Saharan Africa.

A drop in urban childbearing is the principal reason for overall decline in fertility levels. Between the 1992 and 2001-2002 surveys, the urban TFR fell by 1.5 births, from a rate of 5.8 to 4.3 births per woman. In contrast, the rural TFR remained essentially stable over this period (7.1 and 6.9 births per woman, respectively). Women who have at least some secondary education experienced a steady decline in fertility, with the TFR for such women dropping by one birth over the period covered by the three ZDHS surveys.

The length of the interval between births influences the overall fertility level, as well as the health status of the mother and child. There is now a new international consensus that the optimal interval between births is at least 36 months. In Zambia, 41 percent of births occur at least 36 months after a previous birth. Although 16 percent of births occur less than 24 months after a previous birth, this is an improvement from 1996 when almost one in five births (19 percent) occurred less than two years after a previous birth. The median birth interval is 33 months, 3 months shorter than the "safe" birth interval. This represents a one-month increase from the 1996 ZDHS median birth interval (32 months).

Childbearing begins early in Zambia. By age 18 almost half of women age 45-49 have had their first birth. Most women typically become mothers before the age of 20, with 60 to 70 percent of women in all age groups having given birth by that age. The median age at first birth for women age 25-49 is 18.7 years. Three in ten teenage women (age 15-19) have begun childbearing, with 26 percent having had a child and 6 percent pregnant with their first child.

A comparison of the data from the 1996 and 2001-2002 ZDHS surveys indicates that there has been an increase in the median duration of postpartum amenorrhoea from 11.5 months to 13.3 months. There has been little change in the median duration of sexual abstinence or insusceptibility to the risk of pregnancy following childbirth.

FAMILY PLANNING

Knowledge of contraceptive methods is almost universal in Zambia, with 98 percent of all women and men knowing at least one method of family planning. The male condom and the pill are the most widely known methods among both women and men. Knowledge of family planning methods has increased steadily over the last decade. For example, knowledge of at least one contraceptive method has increased among women from 89 percent in 1992 to 96 percent in 1996 to the current level of 98 percent. Since 1996, the mean number of family planning methods known has increased from 5.3 to 6.8 for women and from 5.0 to 6.1 for men.

Mass media are important sources of information on family planning. Almost half of women (49 percent) and six in ten men (62 percent) have heard or seen a family planning message on the radio, television or in a newspaper/magazine. Radio is the most frequent source of family planning messages for both women (46 percent) and men (57 percent).

The 2001-2002 ZDHS data indicate that 70 percent of currently married women and 81 percent of currently married men have used a family planning method at least once in their lifetime. There is a noticeable discrepancy between ever use and current use, with slightly more than one-third of currently married women (34 percent) currently using some method of contraception. Modern methods of contraception are more commonly used than traditional methods; almost one-fourth of currently married women use modern methods (23 percent) compared with about one in ten who use traditional methods (12 percent). The pill is the most widely used modern method (12 percent), while withdrawal is the most popular traditional method (5 percent).

Six in ten women currently using modern methods obtain their method from the public sector (61 percent), 20 percent from the private medical sector, and 17 percent from other sources such as shops and community-based agents.

Contraceptive use in Zambia has increased over the past decade from 15 percent in

1992 to 26 percent in 1996 and 34 percent in 2001-2002. Considering specific methods, the largest gains in use during the ten-year period are observed for contraceptive pills and injectables.

The large majority of currently married women who know of family planning approve of its use (87 percent). Moreover, two-thirds of married women who know of a contraception method believe their husband approves of family planning.

The majority of currently married women (71 percent) and currently married men (58 percent) either want to space their next child or want no more children, and thus have a potential need for family planning. In Zambia, almost one in three currently married women has an unmet need for family planning (27 percent) – 17 percent for spacing their next birth and 11 percent for limiting births. If the unmet need for family planning of all currently married women who say they want to space or limit their births were met, the contraceptive prevalence rate in Zambia would increase from 34 to 62 percent.

MATERNAL HEALTH

Use of professional maternity care is common in Zambia: more than nine in ten mothers who had a live birth in the five years preceding the survey received antenatal care from a health professional; only 2 percent received antenatal care from a traditional birth attendant or other person. Almost three-quarters of these women had four or more antenatal care visits during their pregnancy (72 percent).

Among mothers who received antenatal care, over three-fourths were given at least one tetanus toxoid injection during pregnancy for their most recent birth (75 percent). Over the past 10 years, there has been a decrease in the proportion of women who received at least one tetanus toxoid injection during their most recent pregnancy, from 81 percent in 1992 and 85 percent in 1996 to the current level of 75 percent. The 2001-2002 ZDHS data show that seven in ten mothers receive iron tablets, syrup or folic acid during pregnancy. Ninety-four percent of mothers have their height measured during antenatal care visits, while 25 percent are weighed. Eighty-seven percent of mothers had their blood pressure measured during their antenatal care, and urine and blood sampling was done for 25 and 44 percent of mothers, respectively.

The 2001-2002 ZDHS results show that slightly more than four in ten births are delivered in a health facility. A similar proportion are delivered by a health professional: 39 percent by a nurse/midwife and the remaining 5 percent by a doctor or clinical officer. Almost four in ten births are delivered with assistance from a relative or friend. A comparison between the three ZDHS surveys shows a steady decline in the proportion of births that are delivered in a health facility, from 51 percent in 1992 and 47 percent in 1996 to the current level of 44 percent.

Postnatal care is not common in Zambia. Less than one in four women who deliver outside a health facility receive postnatal care (23 percent).

The 2001-2002 ZDHS collected data allowing measurement of maternal mortality. The maternal mortality ratio during the seven-year period prior to the 2001-2002 ZDHS was estimated at 729 maternal deaths per 100,000 live births. This represents a rise from 649 maternal deaths per 1000,000 live births estimated from the 1996 ZDHS.

CHILD HEALTH

At current mortality levels, one in six Zambian children die before the fifth birthday, (under-five mortality rate of 168 deaths per 1,000 birth), with slightly more than half of these deaths occurring during the first year of life (infant mortality rate of 95 deaths per 1,000 births). A comparison of the three ZDHS surveys, however, shows that mortality among young children has declined from the fairly stable levels observed in the late 1980s and early to mid-1990s. Infant mortality, which had shown a modest increase from 107 to 109 deaths per 1,000 births between the 1992 and 1996 surveys, fell to 95 deaths per 1,000 births in the 2001-2002 ZDHS. Under-five mortality is 15 percent lower now than it was five to nine years ago, with the pace of decline very similar to the decline in infant mortality over the same time period (13 percent).

Early childhood mortality is consistently lower in urban areas than in rural areas. Maternal education is strongly correlated with childhood mortality. Infant mortality is 32 percent

lower and under-five mortality 48 percent lower among mothers with some secondary education than among uneducated mothers.

Survival of children is strongly associated with proper immunisation and treatment of childhood illnesses. According to the World Health Organization, a child is considered fully vaccinated if he or she has received a BCG vaccine, three doses of DPT vaccine, at least three doses of polio vaccine, and one dose of measles vaccine. Only 57 percent of Zambian children are fully vaccinated by 12 months of age; however, 70 percent of children 12-23 months are fully vaccinated, with only 3 percent of children not having received any vaccine.

Looking at coverage for specific vaccines, 94 percent of children 12-23 months received the BCG vaccine, 84 percent received measles vaccine, and 80 percent received the recommended three doses of DPT and polio vaccines.

Comparing the three ZDHS surveys, there was a substantial rise in vaccination coverage in the early to mid-1990s. In contrast, between 1996 and 2001-2002, there was an 11 percent decline in the proportion of children 12-23 months who are fully immunised.

Vitamin A is a micronutrient found in very small quantities in some foods. It is considered essential for normal sight, growth, and development. Zambia has recently introduced a programme of vitamin A supplementation for children 6 to 72 months through health services and community campaigns. Data from the 2001-2002 ZDHS show that two-thirds of children 6-59 months are reported to have received a vitamin A supplement in the previous 6 months.

Fifteen percent of children under five years of age had symptoms of acute respiratory infection (ARI) in the two weeks preceding the survey. Seven out of ten children with ARI symptoms were taken to a health facility or provider for treatment.

Forty-three percent of children under five were reported to have been ill with fever and/or convulsions in the two-week period preceding the survey. Among these children, more than half took antimalarial drugs (52 percent) and almost four in ten took antimalarial drugs the same or next day (37 percent).

According to mothers' reports, around 20 percent of children under age five had diarrhoea at some time in the two weeks before the survey. Slightly more than four in ten children who were ill with diarrhoea were taken to a health facility (43 percent). Mothers reported that almost two-thirds of children with diarrhoea (67 percent) were treated with some form of oral rehydration therapy; more than half (53 percent) were given a solution prepared with oral rehydration salts (ORS); and four in ten were given increased fluids. Twenty-one percent of children with diarrhoea did not receive any type of treatment.

BREASTFEEDING AND NUTRITION

Breastfeeding is nearly universal in Zambia: 98 percent of children born in the five years preceding the survey were breastfed. More than half of infants are put to breast within an hour of delivery and 90 percent are breastfed within the first day. The median duration of breastfeeding is 21 months, a slight increase from 1996 (20 months).

Despite the high prevalence of breastfeeding in Zambia, the majority of infants are not exclusively breastfed for the first six months of life. Only four in ten infants under 6 months of age are exclusively breastfed in Zambia. Complementary feedings starts early; more than one-third of children under 6 months of age are given complementary foods.

According to the 2001-2002 ZDHS findings, the level of malnutrition among children under five is significant, with almost half of Zambian children (47 percent) stunted (short for their age), 5 percent wasted (thin for their age), and more than one-fourth underweight (28 percent). Generally, children who live in rural areas and children of uneducated mothers are more likely to be malnourished than other children.

The nutritional status of women is another issue of importance. The mean height of Zambian women is 158 cm. Only 2 percent of women are shorter than the critical height of 145 cm. Fifteen percent of women fall below the cut-off of 18.5 for the body mass index (BMI) - an indicator used to measure the level of chronic energy deficiency among adults. In general, very

young women (age 15-19) and rural women are more likely than other women to suffer from chronic energy deficiency.

WOMEN'S STATUS AND VIOLENCE AGAINST WOMEN

The 2001-2002 ZDHS provides information on the status of women in Zambia, and on physical and sexual violence against women. Overall, more than one in ten women age 15-49 have no education (12 percent) and women are generally less educated than men. Four in ten women in Zambia are illiterate.

More than half of women were working at the time of the survey (55 percent). Most women work seasonally (53 percent). Agriculture is the predominant sector of the economy, employing 54 percent of women in the 12 months preceding the survey. Forty-two percent of all working women in Zambia are either paid in kind or not paid at all. Women working in the non-agricultural sector are more likely to earn cash than women working in agriculture. Among currently married women who earn cash for their work, 41 percent report that they alone make decisions about how their earnings will be used and 32 percent report that they decide jointly with their husband.

Women in the 2001-2002 ZDHS were asked about their beliefs on wife-beating. A large majority of women (85 percent) believe that a husband is justified in beating his wife for at least one reason. Almost eight in ten women believe that a husband is justified in beating his wife if she goes out with another man. A slightly smaller proportion agree that if a woman neglects her children (61 percent), or argues with her husband (52 percent), then he is justified in beating her.

The 2001-2002 ZDHS found that more than half of women report having experienced beatings or physical mistreatment since the age of 15, and almost one in four women (24 percent) experienced physical violence in the 12 months preceding the survey. Among physically abused women currently in union, almost eight in ten report their current husband/partner as a perpetrator of the violence, while among never-married women who experienced physical abuse, the mother or father is the most commonly reported perpetrator (35 percent).

Overall, 15 percent of women report having experienced sexual violence by a man and 8 percent reported such experience in the 12 months preceding the survey. Among ever-married women who ever experienced sexual violence, the current husband/partner is reportedly the most common perpetrator of such violence (37 percent). More than four in ten never-married women report their current boyfriend as the perpetrator of sexual violence.

HIV/AIDS AND STI-RELATED KNOWLEDGE AND BEHAVIOUR

General awareness of HIV/AIDS is nearly universal among men and women of reproductive age in Zambia. A large majority of respondents (79 percent of women and 76 percent of men) know someone personally who has HIV/AIDS or has died of AIDS.

Given the high levels of HIV/AIDS awareness, it is not surprising that 78 percent of women and 86 percent of men know two or more effective ways to avoid HIV infection. More specifically, 72 percent of women and 79 percent of men mention the use of the condom as a specific way to avoid HIV infection, while 82 percent of women and 86 percent of men mention limiting the number of sexual partners/staying faithful to one partner – all of which are considered to be programmatically important ways of avoiding HIV/AIDS. Respondents who live in urban areas and those with more education are more likely to know about HIV/AIDS and ways to avoid getting infected with HIV.

Knowledge of ways that HIV can be transmitted is important in preventing the spread of the disease in a population. Most women recognize that the HIV virus can be transmitted from a mother to a child during pregnancy (79 percent), during delivery (65 percent), and by breastfeeding (71 percent). Almost eight in ten women and men know that a healthy-looking person can have the AIDS virus.

Overall, only 9 percent of women and 14 percent of men have been tested for HIV. The more educated women and men and those living in urban areas are more likely to have been tested for HIV than other respondents. Roughly

two-thirds of women and men who have not been tested for AIDS say they want to be tested.

One in ten women and men do not know of any sexually transmitted infections (STIs) other than HIV. Only 3 percent of women and 5 percent of men who have ever had sex reported having an STI or symptoms associated with an STI in the 12 months before the survey. More than half of women and men who had an STI in the 12 months preceding the survey sought advice or treatment from a clinic, hospital, or private doctor. Twenty-two percent of women and 32 percent of men with an STI or associated symptoms did not inform their partner, and one in three women and men with an STI took no action to protect their partner.

SYPHILIS AND HIV TESTING

The 2001-2002 ZDHS was the first nationally representative survey in Zambia to include voluntary syphilis and HIV testing. Syphilis test results were linked to the ZDHS interview data with precautions taken to ensure confidentiality of the respondents. HIV testing was carried out anonymously and the test results could not be linked to the respondent's individual information, except for age, sex, residence (urban-rural), and province.

Overall, 6 percent of women and 8 percent of men age 15-49 in Zambia tested positive for syphilis. The syphilis prevalence rate is slightly higher among urban residents of Copperbelt and Lusaka than among those living in rural areas and other provinces. Sixty-three percent of men and 28 percent of women who were found to be syphilis positive have used a condom at some time in their life.

The 2001-2002 ZDHS data on HIV testing found that of the individuals tested, 16 percent were HIV positive. Women are more likely to be HIV-positive than men (18 percent and 13 percent, respectively). Overall, the proportion HIV-positive rises with age from a level of 5 percent among respondents 15-19 to 25 percent among those 30-34, before dropping to 17 percent among those age 45-49. HIV prevalence is more than twice as high in urban areas as in rural areas (23 percent and 11 percent, respectively).

ZAMBIA

